

Association between Shifting Work and Musculoskeletal Disorders among Workers Working in Steel Industry Application of Multivariate Logistic Regression

#### ARTICLE INFO

Article Type Original Research

**Authors** Vakili-Basir A. <sup>1</sup> PhD candidate Gholami-Fesharaki M.<sup>1</sup> PhD

#### How to cite this article

Vakili-Basir A, Gholami-Fesharaki M. Association between Shifting Work and Musculoskeletal Disorders among Workers Working in Steel Industry . International Journal of Musculoskeletal Pain Prevention. 2018;3(4): 107-113.

<sup>1</sup>Biostatistics Department, Faculty of Medical Sciences, Tarbiat Modares University, Tehran, Iran.

\* Correspondence

Address: Department of Biostatistics, Faculty of Medical Sciences, Tarbiat Modares University, Ale-Ahmad Highway, Tehran, Iran. P.O. Box: 14115-111 Phone: +98 (21) 82883578 Fax: +98 (21) 82883578

#### Article History Received: April 20, 2019

Accepted: May 5, 2019 ePublished: May, 5, 2019

#### ABSTRACT

**Aims:** Due to the increasing development of industries and the need for continuous and intensive work, work-related diseases, such as Work- Related Musculoskeletal Disorders (WRMSD), have increased. The present study was carried out to evaluate the association between shifting work and WRMSD.

**Instruments &Methods:** This cross-sectional study was carried out on 300 male workers recruited from Mobarakeh Steel Company Esfahan, Iran during May 2015 till June 2016. The participants in this study were selected by two-stage random sampling (stratify-cluster sampling) in 3 shift categories (day worker, ordinary and rotation shifting worker) and then responded to the Nordic Musculoskeletal Questionnaire (NMQ). Data were analysed using SPSS software (version 24) and ML-win software (version 2.3).

**Findings:** The mean age of the participant was 42.4(SD=8.19), 39.82(SD=8.48) and 44.4(SD=3.57) years in day worker, ordinary and rotation shift worker respectively. The results showed that over the past seven days, the risk of shoulder disorders in ordinary shifts was significantly (OR=1.66, P-value=0.04) higher than the rotational shift. The risk of elbow disorders during the seven days or the past 12 months in ordinary shifts was significantly (OR = 0.38, P-value=0.01) lower than the rotational shift worker.

**Conclusion:** According to the result of this study, considering shifting work greatly is helpful in preventing WRMSD. Therefore, it is necessary to pay more attention to working conditions and risk factors for shoulder, elbow, and legs disorders.

Keywords: Musculoskeletal Disorders, Shifting Work, Steel Industry.

### Introduction

Today's industrial world has brought the growing trend of production rate in shorter time. Therefore, it is not possible such as speed and production without shifting work. In general, shifting work is work that takes place on a schedule outside the traditional time means from 9 am – 5 pm daily <sup>[1]</sup>. In different jobs, depending on the type of work and the need for more workforces, the shifting work has been defined accordingly. The previous study showed that shifting worker tends to health problems like overweight <sup>[2-4]</sup>, high-blood pressure <sup>[5, 6]</sup>, high lipid disorder<sup>[7]</sup>, sleep disorders<sup>[8]</sup> and WRMSDs <sup>[9, 10]</sup>. The previous

study also showed that working in unusual shifts reduces awareness and attention during the day <sup>[11]</sup>. On the other hand, reduced attention and vigilance during daily activities could cause a lot of Musculoskeletal Disorders (MSD) and financial and psychological losses <sup>[12]</sup>. Now day WRMSDs in the industries has been continuously increased [13-<sup>16]</sup>. WRMSDs are one of the most common causes of occupational injuries in developing countries which refers to any tissue damage to the musculoskeletal and nervous system that affects the individuals' function<sup>[17]</sup>. Furthermore, they lead to direct and indirect costs and high [18work-related absenteeism

\**Corresponding Author:* Department of Biostatistics, Faculty of Medical Sciences, Tarbiat Modares University, Ale-Ahmad Highway, Tehran, Iran Postal Code: 14115-111 Tel: +98 (21) 82883578 Fax: +98 (21) 82883578 Mohammad.gholami@modares.ac.ir

#### Association Between Shifting Work and ...

<sup>20]</sup>. WRMSDs, in addition to ergonomic exposures, depends on other factors such as age, sex, body Mass Index (BMI), daytime work, mental stress, and physical fitness <sup>[21-</sup> <sup>25]</sup>. In other hands, in different jobs, type of working and the need for more workforce, working in unallowable and unusual hours of the day could be resulted in WRMSDs. Regarding the importance of social and sanitary effects of -shifting work and the occurrence of muscular disorders, especially in night shifts, and the association of some of these disorders with shifting work and its related outcomes, this study aimed to assess the association between shifting work and WRMSDs among steel company workers.

#### **Instruments and Methods**

This cross-sectional study conducted in Esfahan Mobarakeh Steel Company (EMSC), Iran from May 2015 to June 2016. The aim of this study was to investigate the effects of shifting work on WRMSDs in all staff and male workers who work in the Company. This study was carried out on a sample of 300 people, using a two-stage (stratifiedcluster) sampling method. In this study, we use the formula  $(n = (Z_{\underline{\alpha}} + Z_{\beta})^2 / d^2 + 1)$  for estimating the sample size by considering  $\alpha = 0.05$ ,  $\beta = 0.1$  and d = 0.2. The sample size was calculated as 263 workers. Considering 14% of the sample loss, 37 samples were added and finally, 300 workers considered to be studied in this study. In this study, the willingness to participate, formal or contractual employment and having at least two years of work experience at EMSC considered as inclusions criteria Moreover, unwillingness to participate, expulsion, death or retirement during study considered as exclusions criteria. The participants were fully satisfied and confidentially information was considered with researchers in this study. The Medical Ethics Committee of the Tarbiat Modares University (TMU) approved the ethical protocol of this study (IR.TMU.REC.1396.589). In this study, the definition of shifting schedule fully described by Gholami et al paper<sup>[18]</sup>. In this study, participate fill out demographical questioner addition in Nordic Questionnaire Musculoskeletal (NMQ). The NMQ is yes/no structured interview questionnaire includes the human body image that is divided into nine parts of body including neck, shoulder, elbows, wrists and hands, waist up, lower back, thighs, knees, ankles, and legs. In this questionnaire, participants provided information on their musculoskeletal disorders in the past 12 months (outbreak) and 7 days ago (starting point) <sup>[12]</sup>. The validity and reliability of this questionnaire reported in previous the studies [18, 26].

#### **Statistical Analysis**

In this study, we used multilevel logistic modelling for analysis multivariate logistic regression. In the first multivariate binary, the responses were restructured to one column and each response considered as a repeat of one response. Actually, this approach tends to share parameter in joint modelling <sup>[18]</sup>. Since multilevel models consider the correlation between observations, they provide more accurate results than compared to simple logistic regression analysis [27, <sup>28]</sup>. In multivariate responses, there is the possibility of data loss which can lead to loss of information and missed values. The presence of missed values may decline the accuracy of calculated statistical indicators and subsequently increase the complexity of the model and statistical methods. Since results from incomplete data can be resulted in bias, these data analysis should be done in inappropriate way<sup>[11, 29-31]</sup>. Multilevel model is a suitable method because it does not require data with balanced structure [27, 29, 32]. Data were analysed by ML-win and SPSS software. Descriptive statistics (mean, standard

Downloaded from ijmpp.modares.ac.ir on 2024-05-20

deviation, frequency, and percentage) were used to describe the demographic characteristics. The adjusted OR and Mann–Whitney index were used for testing the hypotheses. Normality of the data was checked by using the Kolmogorov-Smirnov test. The significance level of less than 5% was used as a meaningful level of acceptable.

## **Findings**

The demographical information of the participants according to the shifting schedule are presented in Table (1) and Table (2). According to the results of these two tables, we can see rotation shifting workers are older, more experienced, more smoker, and lower educated and lower weighted rather than day worker and ordinary shifting workers. Since age, experience, education and smoke status were significantly different, so in final analysis such variable was controlled. The results of the Odds Ratio (OR) are presented in Table (3). As it is shown in this table, the risk of MSD in rotation shifting is greater than day shifting worker. It can also be seen that the risk of shoulder disorders in the ordinary shift was higher (OR=1.66) than day shifting worker significantly over the past seven days (P=0.04). The risk of elbow disorders during the seven days or the past 12 months in ordinary shifting workers was reported 62% (OR=0.38) that was less than day shifting worker (P=0.01). Other disorders were not significant different according to the shifting schedule.

# Discussion

Because of the importance of WRMSDs and shifting work, this study was performed to explore the association between shifting work and WRMSDs among steel company workers. The result of this study showed that positive risk of WRMSDs in rotation shift rather than day shifting worker. This finding was in the line with the previous study <sup>[33, 34]</sup> that can be attributed to shifting work experience. Jeff et al reported that diseases such as high blood pressure, gastrointestinal ulcers, chronic worries, asthma and allergies, WRMSDs, and severe obesity among middle-experience shifting workers have been around 40% more than shifting workers with less experience <sup>[35]</sup>. Thus, it can attribute to the high work pressure in shifting work. Choobineh et al showed that significant signs of WRMADs are high in nurses with high physical pressures. Back pain is the most prevalent problems among such nurses [36]. The results also showed the decreasing effect of working in shifting work on risk of elbow disorders. Such phenomena can be explaining by a higher risk of knee disorder in day shifting worker because of their job position rather than shifting worker<sup>[37]</sup>. In contrast with Lipscomb et al study <sup>[9]</sup> and Adel et al. the present study have not shown any significant relationship between shifting work and neck, wrists/ hands, upper and lower back, one/both hips/ thighs/buttocks, one/both knees , one/both ankles/feet disorders.

# **Strength and Limitation**

A limitation of this study is sample selection from a single company and job category. In contrast, performing of risk assessment for all staffs without considering their knowledge, selecting large sample and use of reliable statistical analysis methods were considered as the strength points of this study.

# Conclusion

According the findings of this study, it could be concluded that early preventions such as the replacement of mechanized methods instead manual methods, ergonomic modification of the work environment, the training of the correct use of work tools and correct posture when using these tools are effective in preventing WRMSDs in shifting worker. It is also necessary to pay more attention to working conditions and risk factors for shoulder, elbow, and legs.

		shift schedule						_
Variables	Measurements	Day shifting		Ordinary shifting		Rotation shifting		-
		Mean	SD	Mean	SD	Mean	SD	P-value
Age	year	42.40	8.19	39.82	8.48	44.40	3.57	0.005
Experience	year	16.01	7.99	15.17	7.67	20.48	2.74	0.006
BMI*	Kg/m2	27.08	4.04	26.08	3.13	25.52	3.38	0.057

<b>Table 1</b> Quantitative demographic variables according to the groups of schedules of work shi	ifting
--	--------

<sup>\*</sup>BMI: Body Mass Index

Table 2: Qualitative demographic variables according to the groups of schedules of work shifts

		shift schedule						
Variable	level	Day s	hifting	Ordina	ry shifting	Rota	tion shifting	Darahaa
		Ν	%	Ν	%	N	%	P-value
Education	Diploma or Lower	82	59.0	124	91.2	25	100.0	<0.001
	Upper Diploma	57	41.0	12	8.8	0	0.0%	
Marriage	Married	131	94.2	123	90.4	23	92.0	0.494
	Single	8	5.8	13	9.6	2	8.0	
Smoking	No	46	90.2	58	71.6	8	47.1	<0.001
	Yes	5	9.8	23	28.4	9	52.9	

# Acknowledgments

The authors acknowledge research deputy of TMU for financial support. The authors wish to thanks all the personnel, especially the staff of industrial medicine of EMSC, for their cooperation throughout the study.

### **Ethical permission**

Participants in the study, after hearing the explanations of the researchers, read the consent form and signed it. Then they entered into the study. Medical Ethics Committee of faculty of medical sciences of TMU approved the

# study (code number: IR.TMU.REC.1395.398). Competing interests

There are no conflicts of interest for this study.

# Authors' contributions

V.B. Collected the data analysed and prepared the first draft of the article. Gh. F.M supervised all parts of the project.

### **Funding / Support**

This project was funded by Tarbiat Modares University (Grant number: 52D/250, 2018/04/16).

Disorders		Recent 7 Days	Recent 12 Months	Recent 7 Days or 12 Months				
	Day shifting	Considered as reference category						
Nl-	Ordinary Shifting	1.16 (0.51,2.64)	1.16 (0.51,2.64) 1.31 (0.66,2.6)					
Neck	Rotation shifting	2.29 (0.72,7.33)	2.94 (1.01,8.54)	3.44 (1.17,10.13)				
	P-value	0.37	0.14	0.08				
	Day shifting	Considered as reference category						
Charaldana	Ordinary shifting	1.66*(0.48,5.76)	1.35 (0.52,3.48)	1.35 (0.52,3.48)				
Shoulders	Rotation shifting	6.06 (1.43,25.67)	2.84 (0.8,10.09)	2.84 (0.8,10.09)				
	P-value	0.04	0.27	0.27				
	Day shifting	Considered as reference category						
E11	Ordinary shifting	0.52 (0.08,3.21) 0.15 (0.02,1.31)		0.38**(0.09,1.57)				
EIDOWS	Rotation shifting	4.27 (0.78,23.27)	2.48 (0.53,11.55)	3.91 (1.03,14.77)				
	P-value	0.07	0.06	0.01				
	Day shifting	Considered as reference category						
	Ordinary shifting	1.92 (0.86,4.29)	1.16 (0.54,2.46)	1.55 (0.74,3.25)				
wrists/Hands	Rotation shifting	2.55 (0.79,8.23)	1.73 (0.56,5.4)	2.21 (0.73,6.68)				
	P-value	0.18	0.64	0.3				
	Day shifting	Considered as reference category						
User en De els	Ordinary shifting	1.13 (0.34,3.73)	1.43 (0.53,3.86)	1.71 (0.65,4.52)				
Upper Back	Rotation shifting	1.55 (0.27,8.74)	1.71 (0.4,7.43)	1.71 (0.4,7.43)				
	P-value	0.88	0.7	0.53				
	Day shifting	Considered as reference category						
	Ordinary shifting	0.89 (0.44,1.8)	1.18 (0.62,2.24)	1.23 (0.65,2.35)				
Lower Back	Rotation shifting	1.88 (0.65,5.46)	3.21 (1.03,10.01)	3.21 (1.03,10.01)				
	P-value	0.37	0.13	0.13				
	Day shifting	Considered as reference category						
One/both Hips/	Ordinary shifting	1.35 (0.31,5.87)	1.35 (0.31,5.87)	1.21 (0.33,4.48)				
Thighs/Buttocks	Rotation shifting	4.27 (0.78,23.27)	6.1 (1.22,30.33)	4.5 (1,20.21)				
	P-value	0.21	0.05	0.1				
	Day shifting	Considered as reference category						
	Ordinary shifting	1.08 (0.51,2.32)	1.24 (0.62,2.47)	1.12 (0.57,2.19)				
Une/both Knees	Rotation Shift	2.77 (0.93,8.27)	1.5 (0.51,4.41)	1.91 (0.67,5.48)				
	P-value	0.16	0.72	0.48				
	Day shifting	Considered as reference	e category					
One/both Ankles/	Ordinary shifting	0.92 (0.39,2.14)	0.92 (0.39,2.14) 0.69 (0.31,1.56)					
Feet	Rotation shifting	1.76 (0.53,5.89)	1.33 (0.41,4.34)	1.59 (0.52,4.93)				
	P-value	0.55	0.48	0.4				

**Table 3:** The odds ratio of musculoskeletal disorders in ordinary and rotation shifting worker compared with day shifting worker.

## References

- 1. Gholami-Fesharaki M, Kazemnejad A, Zayeri F, Rowzati M, Sanati J, Akbari H. Multicenter historical cohort study of the relationship between shift work and blood pressure. ARYA Atheroscler (2014);10 (6):287-292:PMC4354080.
- Buchvold HV, Pallesen S, Waage S, Bjorvatn B. Shift 2. work schedule and night work load: Effects on body mass index-a four-year longitudinal study. Scand J Work Environ Health (2018);44 (3):251-257.
- Buden JC, Dugan AG, Faghri PD, Huedo-Medina TB, 3. Namazi S, Cherniack MG. Associations among Work and Family Health Climate, Health Behaviors, Work Schedule and Body Weight. Journal of occupational environmental medicine (2017);59 (6):588-99: 10.1097/JOM.000000001049:PMC5477673.
- Gholami Fesharaki M, Kazemnejad A, Zayeri F, 4. Rowzati M, Akbari H. Relationship between shift work and obesity a retrospective cohort study. Journal Mil Med (2012);14 (2):93-7.
- 5. Bernardes Souza B, Mussi Monteze N, Pereira de Oliveira FL, Magalhães de Oliveira J, Nascimento de Freitas S, Marques do Nascimento Neto R, et al. Lifetime shift work exposure: association with anthropometry, body composition, blood pressure, glucose and heart rate variability. Journal of occupational environmental medicine (2015);73 (3):208:1011360emed-2014-10429.
- Su T-C, Lin L-Y, Baker D, Schnall PL, Chen M-F, Hwang 6. W-C, et al. Elevated blood pressure, decreased heart rate variability and incomplete blood pressure recovery after a 12-hour night shift work. Journal of occupational environmental medicine (2008);50 (5):380-6:https://doi.org/10.1539/joh.L7056.
- Akbari H, Mirzaei R, Nasrabadi T, Gholami-7. Fesharaki M. Evaluation of the effect of shift work on serum cholesterol and triglyceride levels. Iranian Red Crescent Medical (2015);17 (1): 10.5812/ ircmj.18723.
- 8. Kuetting DL, Feisst A, Sprinkart AM, Homsi R, Luetkens J, Thomas D, et al. Effects of a 24-hr-shiftrelated short-term sleep deprivation on cardiac function: A cardiac magnetic resonance-based study. Journal of sleep research (2018):e12665: https:// doi.org/10.1111/jsr.12665.
- 9. Lipscomb JA, Trinkoff AM, Geiger-Brown J, Brady B. Work-schedule characteristics and reported musculoskeletal isorders of registered nurses. Scandinavian journal of work, environment health (2002);28 (6):394-401:http://dx.doi.org/10.5271/ sjweh.691.
- 10. Yasobant S, Rajkumar PJIjoo, medicine e. Workrelated musculoskeletal disorders among health care professionals: A cross-sectional assessment of risk factors in a tertiary hospital, India. (2014);18 (2):75-81:10.4103/0019-5278.146896:PMC4280781.
- 11. Heck RH, Thomas SL. An introduction to multilevel modeling techniques: MLM and SEM approaches using Mplus. 3rd Edition ed. New York

International Journal of Musculoskeletal Pain Prevention

(2015): Routledge. 5 March 2015. https://doi. org/10.4324/9781315746494.

- 12. Asghari M, Ahmadnezhad I, Rahmani A, Sadeghi A, Abbassinia M, Rezaie E, et al. Investigation of disorders and problems caused by shift work in an automotive industry. Journal of North Khorasan University of Medical Sciences (2013);5 (1):7-15.
- 13. Marras W. Occupational low back disorder causation and control. Ergonomics (2000);43 (7):880-902:10.1080/001401300409080.
- 14. Mirmohammadi S, Mehrparvar A, Soleimani H, Lotfi MH, Akbari H, Heidari N. Musculoskeletal disorders among video display terminal (VDT) workers comparing with other office workers. Iran Occupational Health (2010);7 (2):11-4.
- 15. Mostaghaci M, Salimi Z, Javaheri M, Hoseininejad S, Salehi M, Davari M, et al. Evaluation of the musculoskeletal disorders and its risk factors in the workers of an agricultural equipmentmanufacturing plant. Occupational medicine Quarterly Journal (2012);3 (3):19-25.
- 16. Smith DR, Leggat PA. Musculoskeletal disorders among rural Australian nursing students. Australian Journal of Rural Health (2004);12 (6):241-5:https:// doi.org/10.1111/j.1440-1854.2004.00620.x.
- 17. Keyserling WM. Workplace risk factors and occupational musculoskeletal disorders, Part 2: A review of biomechanical and psychophysical research on risk factors associated with upper extremity disorders. American Industrial Hygiene Association journal (2000);61 (2):231-43:https:// doi.org/10.1080/15298660008984532.
- 18. Gholami Fesharaki M, Kazemnejad A, Zayeri F, Rowzati M, Akbari H. Historical cohort study of shift work and blood pressure. Occupational medicine (2014);64 (2):109-12:https://doi.org/10.1093/ occmed/kqt156.
- 19. Gonçalves MB, Fischer FM, Lombardi M, Ferreira RM. Work activities of practical nurses and risk factors for the development of musculoskeletal disorders. Journal of human ergology (2001);30 (1-2):369-74:https://doi.org/10.11183/ jhe1972.30.369.
- 20. Nagata T, Mori K, Ohtani M, Nagata M, Kajiki S, Fujino Y, et al. Total Health-Related Costs Due to Absenteeism, Presenteeism, and Medical and Pharmaceutical Expenses in Japanese Employers. Journal of occupational environmental medicine (2018);60 (5):e273-e80:10.1097/ JOM.000000000001291:PMC5959215.
- 21. Cabeças JM. Occupational musculoskeletal disorders in Europe: impact, risk factors and preventive regulations. (2006).
- 22. Hootman JM, Sniezek JE, Helmick CGJJowsh, medicine g-b. Observations from the CDC: Women and Arthritis: Burden, Impact, and Prevention Programs. (2002);11 (5):407-16.
- 23. Koo H-R, Shin Y-S, Chae H-S, Lee K-S. The research of job stress and MSDs symptoms of small

112

plants with agricultural products. Journal of Agricultural Extension Community Development (2011);18 (4):861-77:https://doi.org/10.12653/jecd.2011.18.4.861.

- Mohamadbaigi A, Arjmandzadeh A, Nouri E, Choobineh A. Musculoskeletal disorders among bank computer operators. Salāmat-i kār-i Īrān (2006);3 (2):3-7.
- 25. Peele PB, Xu Y, Colombi A, editors. Medical care and lost work day costs in musculoskeletal disorders: older versus younger workers. International Congress Series Elsevier.; 2005 1280, 214-218.
- Choobineh A, Nouri E, Arjmandzadeh A, Mohamadbaigi A. Musculoskeletal Disorders among Bank Computer Operators. Iranian Journal of Epidemiology (2006);3 (2):3.
- 27. Haji AM, Ayat Elahi SMT, Behboudian J. The Effect of Missing Data in Growth Curves. Journal of Babol University of Medical Sciences (2004);6 (4):23-9.
- Sullivan LM, Dukes KA, Losina E. An introduction to hierarchical linear modelling. Statistics in medicine (1999);18 (7):855-88:https://doi.org/10.1002/ (SICI)1097-0258(19990415)18:7<855::AID-SIM117>3.0.CO;2-7.
- 29. Brady T. West, Kathleen B. Welch, Galecki AT. Linear Mixed Models: A Practical Guide Using Statistical Software. 2nd Edition ed. New York (2014): Chapman and Hall/CRC. https://doi.org/10.1201/ b17198.
- 30. Harvey G. Multilevel Models in Educational and Social Research. (1987).
- 31. McCulloch CE, Searle SR. Generalized, Linear, and Mixed Models (Wiley Series in Probability and

Statistics). taylorfrancis .(2001).1-440. https://www. taylorfrancis.com/books/9780429186561

- 32. Buxton R. Statistics: Multilevel Modeling. Mathematics Learning Support Centre–Technical report.taylorfrancis. (2008).www. statstutor. ac. uk/ resources/uploaded/multilevelmodelling.
- Skela-Savič B, Pesjak K, Hvalič-Touzery S. Low back pain among nurses in Slovenian hospitals: crosssectional study. Int Nurs Rev(2017);64 (4):544-51: https://doi.org/10.1111/inr.12376.
- 34. Wilhelmus Johannes Andreas G, Wernstedt P, Campo M. The impact of a rigorous multiple work shift schedule and day versus night shift work on reaction time and balance performance in female nurses: a repeated measures study. Physiotherapy Theory and Practice (2016);58 (7):737-43:https://doi.org/10.31 09/09593985.2010.481323.
- Wilhelmus Johannes Andreas G, Wernstedt P, Campo M. Work-related musculoskeletal disorders in female Swedish physical therapists with more than 15 years of job experience: Prevalence and associations with work exposures. Physiother Theory Pract (2011);27 (3):213-22:https://doi.org/10.3109/09593985.201 0.481323.
- Choobineh A, Rajaeefard A, Neghab M. Perceived demands and musculoskeletal disorders among hospital nurses. Hakim research journal (2007);10 (2):70-5.
- 37. Zare R, Khazraei T, Choobineh A, Daneshmandi H, Rahimian JT, Rajabi AJSMSJ. Assessment of the Risk of Musculoskeletal Disorders using the Quick Exposure Check Technique among the Workers of a Shipbuilding Company. (2014);2 (4).12-18.