

## Low Back Pain Intensity and Quality of Life among Patients referred to a Pain Clinic in Iran

### ARTICLE INFO

**Article Type**  
Original Research

### Authors

Seyedeh Somayeh Kazemi<sup>1\*</sup> PhD  
Rahman Panahi<sup>2</sup> PhD  
Mohammad Rafighi<sup>3\*</sup> MSc

### How to cite this article

Kazemi SS., Panahi R., Rafighi M. Low Back Pain Intensity and Quality of Life among Patients referred to a Pain Clinic in Iran. IJMPP. 2021; 6(4): 595-600.

<sup>1</sup> Health Network, Disease Unit, Mazandaran University of Medical Sciences, Chaloos, Iran.

<sup>2</sup> Ph.D graduated from Health Education and Promotion, Faculty of Medical Sciences, Tarbiat Modares University, Tehran, Iran.

<sup>3</sup> Department of Education and Human Resources Development, Mazandaran University of Medical Sciences, Sari, Iran.

### \* Correspondence

Address: Department of Education and Human Resources Development, Mazandaran University of Medical Sciences, Sari, Iran.  
Tel: +981133044183  
Fax: +981133352725  
P.O.Box: 4815733971  
Email: mohammadrafigi@yahoo.com

### Article History

Received: June 22, 2021  
Accepted: Sep 13, 2021  
ePublished: Oct 9, 2021

### ABSTRACT

**Aims:** Low Back Pain (LBP) is a major contributor to disability worldwide which affected the Quality of Life (QOL). This study aimed to determine the correlation between pain intensity and QOL in patients with chronic LBP.

**Method and Materials:** In this study, 31 patients with LBP who referred to a pain clinic in Sari, the center city of Mazandaran province of Iran were selected through simple random sampling based on inclusion/exclusion criteria. Data collection was done using a demographic questionnaire, Visual Analog Scale (VAS), and Short Form Quality of Life (SF-36). To analyze the data, descriptive / analytical statistics were used through SPSS version 23.

**Findings:** Of the total number of patients, 51.62% were female (n=16), and 48.38% were male (n=15). The majority of the participants (N=22; 70.97%) aged greater than 60 years old. The result showed the majority of the participants (N= 17) of patients (54.85%) had moderate pain. There was a significant correlation between pain intensity and all dimensions of QOL except social functioning.

**Conclusion:** This study revealed that low back pain could decrease physical and mental functioning of the suffered patients. Although, doing more studies with larger sample and objective measurements, applying different approaches to decrease low back pain is guaranteed.

**Keywords:** Low Back Pain, Pain Intensity, Quality of Life, Patient

### Introduction

Low Back Pain (LBP) refers to pain and discomfort localized in the lumbosacral region, with or without radiating leg pain. This pain is between the costal margins and the inferior gluteal folds, which is usually accompanied by movement limitation [1]. In 2013, WHO estimates the prevalence of LBP about 60-70% in industrialized countries [2]. In 2015, the global point prevalence of activity-limiting LBP was 7.3%, implying that 540 million people were affected at any time. Low back pain is now the number one cause of disability globally [3]. It has been estimated that about 70 % of people in developed countries experienced LBP during their lives [4,5]. The largest increases in disability caused

by LBP in the past few decades have occurred in low-income and middle-income countries, including in Asia, Africa, and the Middle East, where health and social systems are poorly equipped [6].

It has been argued that the prevalence of LBP among general population is between 60 and 80% [7,8]. Low back pain is an extremely common symptom experienced by people of all ages [9-11].

Globally, LBP is among the highest contributors to disability and it is ranked as a reason of Years Lived with Disability (YLD) [3, 12]. In other words, LBP is the leading cause of YLD in both developed and developing countries, and sixth in terms of overall disease burden (disability-adjusted life-years) [13,14]. The key contributors

to LBP and disability include comorbidities, social factors, psychological factors, genetic factors, and biophysical factors [11].

The prevention strategies for LBP can be mentioned to exercise, education and self-care, back belt, shoe insoles, ergonomic interventions at the workplace, ergonomic school furniture, yoga, cognitive behavioral therapy, massage, etc [15-17].

It has been demonstrated that LBP is one of the main reasons for seeking medical care and it causes an enormous medical and economic burden on individuals, families, communities, industry, and governments [18]. LBP can also affect physical function and quality of life (QOL) and lead to decreased function and QOL [5,19].

Quality of life is the general well-being of individuals and communities, outlining negative and positive aspects of life. It observes life satisfaction, including all aspects from physical health, family, education, employment, wealth, safety, security to freedom, religious beliefs, and the environment [20]. In general, it can be pointed out that back pain affects all aspects of life such as physical, financial, spiritual, and psychological, which leading to a lowers QOL [21].

Studies discussed that there is correlation between pain intensity and QoL. It means, that the increased intensity of pain could be resulted in less physical function of the individuals and affected on their mental / social and general health [22,23].

In this study we aimed to investigate the relationship between the intensity of pain and QOL in patients with chronic LBP in a pain clinic. The findings of this study could help support existed evidences.

### Method and Materials

The participants were the population suffering from non-specific LBP referred to a pain clinic in Sari the center of Mazandaran

province in north of Iran. Based on the inclusion and exclusion criteria, we screened 65 patients with LBP, of which 34 patients were excluded. The inclusion criteria in this study were as being suffered from LBP (female and male) , referring to pain clinic in Sari from November to December 2019 and agreed to participate in the study by signing the informed consent. The exclusion criteria were suffering from pathological LBP, spondylosis, and other visceral pain.

The sampling technique in this study carried out randomly and based on patients' code until the number of subjects reached 31 patients. Data collection was done using a demographic questionnaire include (age, gender, married status), Visual Analog Scale (VAS), and Short Form Quality of Life (SF-36). Pain intensity measured through interviews using the VAS. The VAS is a well-known measure of pain intensity [24], which has been widely used in different adult populations [25]. It is a continuous scale comprised, usually 10 centimeters (100 mm) in length, anchored by 2 verbal descriptors ranging from none (score of 0) to worse condition (score of 100) [25,26]. The VAS measurements were from painlessness (0-4 mm), mild pain (5-44 mm), moderate pain (45-74 mm), and severe pain (75-100 mm). In this study, we used a 100mm straight line to assess pain intensity utilizing the usual anchors.

Quality of life is a concept related to one's physical health, psychological state, level of independence, social relationships, personal beliefs, and relation to the environment. This variable is measured through interviews using the short form quality of life (SF-36) questionnaire. SF-36 consists of 36 items tapping into eight subscales: Physical Functioning (PF), Role Physical (RP), Bodily Pain (BP), General Health (GH), Social Functioning(SF), Vitality (VI), Role Emotional (RE) and Mental Health(MH). The value obtained is in the range of 0-100%

where the higher the value, the better QOL [27]. The questionnaire has been validated in Iran [28].

Descriptive statistics (i.e., frequency, percentage, mean, and standard deviation) was applied. The normal distribution of numeric variables was assessed with the Shapiro-Wilk test. Pearson’s correlation test on normal distribution value and Spearman’s correlation test on abnormal distribution value were used. The analysis of data was done using IBM SPSS software version 23.0.

**Findings**

Of the total number of patients, 51.62% were female (n=16), and 48.38% were male (n=15). The majority of the participants aged greater than 60 years old in both gender (Table 1).

**Table 1)** General characteristics of the studied participants

Variables	N=31	%
<b>Gender</b>		
Male	13	41.94
Female	18	58.06
<b>Age</b>		
< 60	9	29.03
≥ 60	22	70.97
<b>Married Status</b>		
Single	1	3.23
Married	24	77.42
Divorced	2	6.45
Widow	4	12.9

Table 2 shows the pain characteristics in the VAS value. (mean = 6.71, SD = 2.06). Correlation between pain intensity and QOL demonstrated a correlation between VAS and physical function, limitations due to physical problems, limitations due to emotional problems, vitality, the feeling of pain, general

health, and mental health. But there was no connection between VAS and social function. The correlation between pain intensity and QOL is presented in Table 3.

**Table 2).** Characteristics of pain intensity of the studied participants

Variables	Mean	SD
VAS	6.71	2.06
	<b>N=31</b>	<b>%</b>
<b>Painlessness</b>	0	0
<b>Mild Pain</b>	8	25.8
<b>Moderate Pain</b>	17	54.85
<b>Severe Pain</b>	6	19.35

**Table 3)** Correlation between pain intensity and quality of life in studied participants.

Dimension of Quality of Life	VAS	
	r	P
<b>Physical Functioning</b>	-0.612	<0.001 <sup>a</sup>
<b>Role Physical</b>	-0.861	<0.001 <sup>b</sup>
<b>Bodily Pain</b>	0.483	0.01 <sup>a</sup>
<b>General Health</b>	-0.350	0.030 <sup>a</sup>
<b>Vitality</b>	-0.463	0.006 <sup>a</sup>
<b>Social Functioning</b>	-0.245	0.12 <sup>b</sup>
<b>Role Emotional</b>	-0.436	0.004 <sup>b</sup>
<b>Mental Health</b>	-0.480	0.001 <sup>a</sup>

<sup>a</sup>Pearson’s Correlation Test

<sup>b</sup>Spearman’s Correlation Test

**Discussion**

This study aimed to investigate correlation between pain intensity and QOL in chronic LBP patients referred to pain clinic in Sari. Chronic LBP diagnosis was performed through physical and neurological examination by chiropractor and orthopedist. The most patients were female (58.06%) and aged over than 60 years old (70.97%). Ste-

fane study et al. (2013) showed of 97 LBP patients, 67 women (69.07%) and 30 men (30.93%) were aged more than 60 years old [29]. As well, in similar studies, the proportion of female was higher than male, and most participants were aged over 60 years old [5,30]. These data revealed that LBP is more prevalent among female and elderly people. The results of present study showed 54.85% of patients had moderate pain and 25.8% suffered from mild pain. Previous studies demonstrated an appropriate educational intervention can help patients to promote their preventive health behaviors and decrease their LBP [31-35].

Evidences showed pain intensity affected the QOL and could decrease it [31,36]. In this study, findings extracted from the SF-36 questionnaire displayed that there was a significant correlation between pain intensity and almost all of the QOL dimensions except social functioning. These results are supported by Nasution' study et al. (2018). The difference was that in Nasution' study, pain intensity was associated with all dimensions of QOL except mental health [5]. Shim' study et al. (2014) compared LBP groups with healthy groups and found significant correlation between pain severity and physical function, social function, vitality, bodily pain, general health, limitations due to economic problems, and role physical [37] that supported our findings. While However, Nasution et al. (2018) found a relationship between pain severity and mental health but not significant [5]. The mental health element is a valid assessment of mental status and is useful for screening for psychiatric disorders. The total QoL score in the Zahra' study et al. (2020) was low in 77.5% of the studied sample and there was a highly significant negative correlation between the total QOL score and total back pain of the studied sample. That means LBP continues to be a common disease [21].

Importantly, both physical and mental health is linked with chronic back pain. Back pain is associated with reduced QOL, and a significant relationship with chronic back pain was found among all measures of physical and mental health [38]. Similarly, Demirtas' study (2013) was carried out regarding the effects of LBP on QOL and functional impairment in nurses suffering from LBP [39]. The nurses who had more pain intensity had significantly worse scores of functional disability, general health, physical function, role physical, social function, and bodily pain domains of SF-36 compared to nurses who had less pain intensity.

Since LBP causes many health-related problems affecting on not only the patient's physical state but also all aspects of their lives, such as mood and health-related QOL. Therefore, health-related quality of life is an important outcome of health in different aspects of physical, mental, and social dimensions. In this regard, improving the QOL through education is an essential issue for health educators. A better understanding of the association between back pain and health-related QOL could facilitate the implementation of new intervention approaches to the prevention and treatment of back pain.

Although this study has strength points, there are some limitations which could effect on its findings. One of the limitations of this study was self-reporting of patient's pain perception that has been subjective and might be the obtained results are not appropriate. As well, the small sample size was another limitation should be considered. Therefore, doing further studies about assessing the relationship between pain intensity and QOL are strongly recommended.

### Conclusion

This study revealed that low back pain could decrease physical and mental functioning

of the suffered patients. Although, doing more studies with larger sample and objective measurements, applying different approaches to decrease low back pain is guaranteed.

### Acknowledgments

The authors of this study were grateful to all the respected participants who took part in this study and Pain Clinic of Bu Ali Hospital in Sari.

### Authors' contribution

SSK was the main investigator, collected and analyzed the data, and wrote the first draft of the manuscript. MR contributed to the writing process. RP contributed to provide the final draft of manuscript. All authors read and approved the final manuscript.

### Conflicts of Interests

No conflict of interest has been declared by the authors.

### Ethical permission

The ethics committee approval was obtained from the pain clinic of Bu Ali Hospital, Sari. All participants were satisfied to entered into the study and completed a written consent form.

### Funding/Supports

None.

### References

1. Comachio J, Magalhães M O, de Moura Campos A P, Silva C e, Marques A P. A cross-sectional study of associations between kinesiophobia, pain, disability, and quality of life in patients with chronic low back pain. *Adv. Rheumatol.* 2018; 58(8): doi.org/10.1186/s42358-018-0011-2
2. Ehrlich G E, Low back pain Bull. WHO. 2003; 81(9): 671-6
3. Vos T, Allen C, Arora M, Barber RM, Bhutta ZA, Brown A, et al. Global, regional, and national incidence, prevalence, and years lived with disability for 310 diseases and injuries, 1990-2015: a systematic analysis for the global burden of disease study 2015. *Lancet.* 2016;388(10053):1545-602.
4. Chou R. Low back pain (chronic). *BMJ Clin Evid.* 2010:1116.
5. Nasution I K, Lubis N D A, Amelia S, Hocin K. The correlation of pain intensity and quality of life in chronic LBP patients in Adam Malik general hospital. *IOP Conf. Series: Earth Environ Sci.* 2018; 125(1): doi :10.1088/1755-1315/125/1/012183
6. Hoy DG, Smith E, Cross M, et al. Reflecting on the global burden of musculoskeletal conditions: lessons learnt from the global burden of disease 2010 study and the next steps forward. *Ann Rheum Dis.* 2015; 74(1): 4-7
7. Deyo RA, Weinstein JN. Primary care - low back pain. *N Engl J Med.* 2001; 344(5): 363-70.
8. Jaromi M, Nemeth A, Kranicz J, Laczko T, Betlehem J. Treatment and ergonomics training of work-related lower back pain and body posture problems for nurses. *J Clin Nurs.* 2012; 21(11-12):1776-84.
9. Hoy D, Bain C, Williams G, et al. A systematic review of the global prevalence of low back pain. *Arthritis Rheum.* 2012; 64(6): 2028-37.
10. Kamper SJ, Henschke N, Hestbaek L, Dunn KM, Williams CM. Musculoskeletal pain in children and adolescents. *Braz J Phys Ther.* 2016; 20(3): 275-84.
11. Hartvigsen J, Hancock MJ, Kongsted A, Louw Q, Ferreira ML, Genevay S, et al. Low back pain 1. *Lancet.* 2018; 391(10137): 2356-67.
12. Kassebaum NJ, Arora M, Barber RM, Bhutta ZA, Brown J, Carter A, et al. Global, regional, and national disability-adjusted life-years (DALYs) for 315 diseases and injuries and healthy life expectancy (HALE), 1990-2015: a systematic analysis for the global burden of disease study 2015. *Lancet.* 2016; 388(10053):1603-58.
13. Hoy D, March L, Brooks P, et al. The global burden of low back pain: estimates from the Global Burden of Disease 2010 study. *Ann Rheum Dis.* 2014; 73(6): 968-74.
14. Global Burden of Disease Study 2013 Collaborators. Global, regional, and national incidence, prevalence, and years lived with disability for 301 acute and chronic diseases and injuries in 188 countries, 1990-2013: a systematic analysis for the Global Burden of Disease Study 2013. *Lancet.* 2015; 386: 743-800.
15. Stochkendahl MJ, Kjaer P, Hartvigsen J, et al. National clinical guidelines for non-surgical treatment of patients with recent onset low back pain or lumbar radiculopathy. *Eur Spine J.* 2018; 27: 60-75.
16. Qaseem A, Wilt TJ, McLean RM, Forciea MA, Clinical Guidelines Committee of the American College of Physicians. Noninvasive treatments for acute, subacute, and chronic low back pain: a clinical practice guideline from the American College of Physicians. *Ann Intern Med.* 2017; 166: 514-30.

17. UK National Institute for Health and Care Excellence. Low back pain and sciatica in over 16s: assessment and management. November 2016. <https://www.nice.org.uk/guidance/ng59> (accessed Nov 7, 2017).
18. Furlan AD, Malmivaara A, Chou R, Maher CG, Deyo RA, Schoene M, van Tulder MW. 2015 updated method guideline for systematic reviews in the Cochrane back and neck group. *Spine*. 2016; 40 (21): 1660-73.
19. Kazemi SS, Tavafian SS, Hidarnia A, Montazeri A. Consequences and factors affecting work-related low back pain among nursing professionals: A qualitative study. *Payesh*. 2019; 18(3): 291- 303.
20. Barcaccia, Barbara (4 September 2013). Quality Of Life: Everyone Wants It, But What Is It?. *Forbes/ Education*. Retrieved 10 May 2019.
21. Zahra N, Sheha E, Elsayed H. Low Back Pain, Disability and Quality of Life among Health Care Workers. *IJPRPM*, 2020, 9(2):34-44.
22. Rostad HM, Puts MTE, Cvancarova Småstuen M, Grov EK, Utne I, Halvorsrud L. Associations between Pain and Quality of Life in Severe Dementia: A Norwegian Cross-Sectional Study. *Dement Geriatr Cogn Disord Extra*. 2017; (7):109-21.
23. Relationship between diastasis of the rectus abdominis muscle (DRAM) and musculoskeletal dysfunctions, pain and quality of life: a systematic review. *Physiotherapy*. 2019; 105 (1): 24-34.
24. McCormack H M, David J D L, Sheather S. Clinical applications of visual analogue scales: a critical review. *Psychol. Med*. 1988; 18(4): 1007-19.
25. Hawker G A, Mian S, Kendzerska T, French M. Measures of adult pain: Visual analog scale for pain (vas pain), numeric rating scale for pain (nrs pain), mcgill pain questionnaire (mpq), short-form mcgill pain questionnaire (sf-mpq), chronic pain grade scale (cpgs), short form-36 bodily pain scale (sf-36 bps), and measure of intermittent and constant osteoarthritis pain (icoap). *Arthritis Care Res*. 2011; 63(11): 240-52.
26. Jensen M P, Karoly P, Braver S. The measurement of clinical pain intensity: a comparison of six methods. *Pain*. 1986; 27(1): 117-26.
27. Ware J E, Sherbourne C D. The MOS 36-item short-form health survey (SF-36): I. Conceptual framework and item selection. *Med Care*. 1992; 30(6): 473-83.
28. Montazeri A, Goshtasebi A, Vahdaninia M, Gandek B. The Short Form Health Survey (SF-36): translation and validation study of the Iranian version. *Quality of life research. Qual Life Res*. 2005; 14: 875-82.
29. Stefane T, Santos A M, Marinovic A, and Hortense P. Chronic low back pain: pain intensity, disability and quality of life. *Acta Paul Enferm*. 2013; 26(1): 14-20.
30. Yong S C, et al. How does chronic back pain influence quality of life in Koreans: A cross sectional study. *Asian Spine J*. 2014; 8(3): 346-52.
31. Kazemi SS, Tavafian SS, Hiller CE, Hidarnia A, Montazeri A. The effectiveness of social media and in-person interventions for low back pain conditions in nursing personnel (SMILE). *Nurs. Open*. 2021; 8: 1220-31.
32. Kazemi SS, Maghbouli R, Rafighi M. Effectiveness of the Back School Program in Employees Suffering from Low Back Pain. *IJMPP*. 2020; 5(2): 329-35.
33. Baillie L, Bacon C.J, Hewitt C.M, Moran R.W. Predictors of functional improvement in people with chronic low back pain following a graded Pilates-based exercise programme. *J Bodyw Mov Ther*. 2019; 23 (1): 211-18.
34. Özdiñç Anar S. The effectiveness of home-based exercise programs for low back pain patients. *J Phys Ther Sci*. 2016; 28 (10): 2727-30.
35. Tavafian SS, Jamshidi A, Shahmohammadi Sh. Low Back Pain Educational Programs and Quality of Life in Women Living With Chronic Low Back Pain: A Semi Experimental Study. *Iran J Health Educ Health Promot*. 2014; 2(1): 49-56.
36. Rahimi A, Vazini H, Alhani F, Monireh Anosheh. Relationship between Low Back Pain with Quality of Life, Depression, Anxiety and Stress among Emergency Medical Technicians. *Trauma Mon*. 2015; 20(2): e18686.
37. Shim J, Lee K, Yoon S, Lee C, Doh J, Bae H. Chronic low back pain in young Korean urban males: The lifetime prevalence and its impact on health-related quality of life. *J Korean Neurosurg Soc*. 2014; 56(6): 482-7.
38. Husky MM, Farin FF, Compagnone P, Fermanian C, Kovess-Masfety V. Chronic back pain and its association with quality of life in a large French population survey. *Health Qual Life Outcomes*. 2018; 16(1):26;16(1):195.doi:10.1186/s12955-018-1018-4.
39. Demirtas RN. The effects of low back pain on quality of life and functional disability in nurses with low back pain. *Ann. Rheum. Dis*. 2013; 71(3):758.