

Predicting Pain Anxiety Symptoms based on Pain Perception with the mediating role of Mental Pain in Musculoskeletal Patients

ARTICLEINFO

Article Type Original Article

Authors

Fatteme Raiisi^{1*} *PhD* Zahra Reyhaninejad kafi² *PhD* Razieh Rahmani³ *Ph.D candidate*

How to cite this article

Raiisi F., Reyhaninejad kafi Z., Rahmani R. Predicting Pain Anxiety Symptoms based on Pain Perception with the mediating role of Mental Pain in Musculoskeletal Patients. IJMPP. 2022; 7(2): 702-707.

¹ Department of Cognitive Sciences, Institute of Cognitive Science Studies, Tehran, Iran.

²Sport Pathology and Corrective Exercises, Physical Education and Sports Science Faculty, Tehran University, Tehran, Iran.

³General psychology, Refah University, Tehran, Iran.

* Correspondence Address: Department of Cognitive Sciences, Institute of Cognitive Science, Shaid Chamran Bulv, Cognitivescience Buvl.; Edalat Sq, Pardis city phase 4, Tehran, Iran Tel: +982176291130 Fax: +98 21 76291140 P.O.Box: 1658344575 Email: Elhamaraiisi@gmail.com

Article History

Received: Feb 11, 2022 Accepted: Feb 28, 2022 ePublished: Apr 20, 2022

A B S T R A C T

Aims: The purpose of this study is to investigate the prediction of pain anxiety symptoms based on pain perception with the mediating role of mental pain in in patients suffering from musculoskeletal disorders.

Method and Materials: This research is a cross-sectional study in which descriptivecorrelational method was applied. The study population was all musculoskeletal patients in Tehran, Iran in 2021. In present study, 300 musculoskeletal patients were selected via candidate sampling method. They completed the Orbach & Mikulincer Mental Pain Scale (2018), Melzack McGill Pain Questionnaire (1975) and Paknejad et al. Pain Anxiety Symptoms (2014). Pearson correlation & bootstrap test were used to analyze data through SPSS-22.

Findings: There was significant negative relationship between pain perception and mental pain and conversely and significant direct positive relationship between pain perception and pain anxiety symptoms (P0.01<). Moreover, itwas shown the mediating role of mental pain in the relationship between pain perception and pain anxiety symptoms in patients with musculoskeletal pain.

Conclusion: To conclude, based on the results, it could be argued that pain is affected by mental components and if the psychological reasons are relieved, the pain will be perceived less severe.

Keywords: Pain Perception, Mental Pain, Pain Anxiety Symptoms, Musculoskeletal Patients.

Introduction

According to International Association for the Study of Pain (IASP), pain is a state that interrupt with daily life and it is described as tissue damage potentially or not^[1]. Pain is expressed in different ways ^[2] and comes in many forms and in any case affects the quality of life of people ^[3]. On the other hand, pain can be associated with various personal components [4] including psychological components as depression, selfefficacy and so on ^[5]. Researches have shown that pain intensity and chronic pain are influenced by psychological and cognitive factors ^[6]. As cited before, pain is reaction of body into illness and injuries. When the causes of pain are disappeared but pain is remained it is called chronic

pain ^[7] and it cause pain anxiety symptoms.

Therefore, many of the factors that perpetuate pain can be psychological^[8]. In psychology, some pain even chronic pain is rooted from our mental states. In another word, mental pain or emotional pain targets our self and its meaning and in the end, distorts our experiences of pain. It accompanies with undesirable feelings ^[9]. Mental pain can be caused by deep depression, which is often caused by depression and can lead to suicide or suicidal ideation ^[10]. According to psychoanalytical approach, mental pain has a secondary benefit for the person to escape from his real self or change it into pain ^[11]. In clinical conditions, researchers have shown; there are reciprocal

Copyright© 2018, TMU Press. This open-access article is published under the terms of the Creative Commons Attribution-NonCommercial 4.0 International License which permits Share (copy and redistribute the material in any medium or format) and Adapt (remix, transform, and build upon the material) under the Attribution-NonCommercial terms

relationships between chronic pain and emotions and its constructs ^[12]. In cognitive neuroscience; emotional or psychological pain can alter pain circuits in our brain and translated it as chronic pain in long time. In another word, neuronal and chemical factors are involved in this phenomenon ^[13].

Pain perception can be facilitated by mental pain and pain anxiety symptoms ^[14]. Chronic pain refers to psychological and environmental factors like anxiety that they can exacerbate conditions for pain. Our different perception of pain is related to genetic, cultural, psychological, environmental factors. Attention, understanding, control and expectations can affect pain perceptions ^[15]. Pain perception is affected by memory, emotional learning and affective component ^[16]. In a study, results showunconscious emotional stimuli of pain can decrease pain perception ^[17]. In another study, researchers found that there is relationship between pain and anxiety stimuli ^[18].

Psychological treatments as Cognitive Behavioral Therapy, improve psychological factors that they are related to chronic pain like; anxiety, depression, psychological problems, risk factors, and low wellbeing^[19]. Physical chronic pain is affected by emotional regulation mechanisms in patients and they are related to their quality of life and life style ^[20]. In cognitive viewpoint, individuals with chronic pain can minimize their pain by attention distortion. In fact, they have strong cognitive mechanisms that they distract themselves from the pain and these mechanisms decrease pain perception in chronic patients ^[21]. A model of pain perception in people with chronic pain syndrome has shown; there are psychosocial variables as pain catastrophic and perceived social support have important roles in in perception of pain and in patients^[22].

The quality of perceived pain can affect the severity of the pain. Hence, it is less common

in the research literature to evaluate the psychological and cognitive aspects of pain. Most previous studies have examined the physical aspects of pain. Few studies have examined the psychological factors of pain in details, such as the relationship between depression and pain ^[5]. Therefore, a study is needed that links the relationship between pain anxiety symptoms and pain perception mediated by mental pain. Because all these variables depend on psychological characteristics. According to these points; the question of this study is; if there is any mediating role for mental pain while relationship between pain perception and pain anxiety symptoms in musculoskeletal patients is assessed?

Method and Materials

This cross-sectional study was conducted through descriptive–correlational method. The study population included all musculoskeletal patients in Tehran, Iran in 2021. In this study, 300 musculoskeletal patients were selected via candidate sampling method. They completed the questionnaires of study online via Instagram and Whatsapp. Then, data were analyzed by SPSS-22. The instruments of this study included as followes:

Demographic questionnaire This scale: included questions regarding age, gender, kind of pain, academic level and so on.

McGill Pain Questionnaire (short form): This scale was provided by Melzack ^[23] and consists of four main dimensions, 20 subgroups and a total of 78 words. In each subgroup the pain intensity increases from top to bottom. Thus, the lowest word has the highest score. The subscales are sensory perception of pain, emotional perception of pain, perception of pain assessment, and various pains. Dworkin et al. reported Cronbach's alpha coefficients for these subscales 0.87, 0.87, 0.83 and 0.86 respectively^[24]. In Iran, a study demonstrated

0.85 for whole questionnaire ^[25].

Mental Pain Scale: This scale was designed by Orbach et al. [26] It has 44 items and 9 subscales as labeled emptiness, irreversibility, emotional flooding, helplessness, lack of control, bewildered, alienation, narcissism/worthlessness and confusion. The rating of this questionnaire is in 5-point Likert. In a study, Guimarães, et al. ^[27] found internal consistency and test-retest reliability and it has concurrent validity with WHO quality of life, Beck depression and Beck anxiety in addicts of Portugal. In Iran, Karami et al. ^[28] demonstrated 6-factor by exploratory factor analysis that these factors provided 0.40.6 % of mental pain variance. Cronbach's alpha for the whole mental pain questionnaire was 0.96.

Pain Anxiety Symptoms Scale: This scale is developed by Paknejad et al. and it has 40 items. Scoring is in the form of 6-point Likert. In Iranian sample, exploratory factor analysis demonstrated 3 factors as avoidance-escape, fear of pain and physical symptoms. The correlation analysis with TSK scale, PCS, scales of DASS-21, pain self-efficacy beliefs and pain behavior questionnaire showed the convergent and divergent validity ^[29].

Findings

In this study, 300 (187 females and 113 males) musculoskeletal patients were participated. Mean age of participants was 31.27 years old (SD=2.81). The mean and standard deviation of patients' responses to research variables are given in Table 1.

Correlational results showed that there was relationship between pain perception and mental pain negatively. Conversely the relationship between pain perception and pain anxiety symptoms was positively and directly significant (P0.01<). In another word, when the pain anxiety symptoms of patients with musculoskeletal pain increases, their pain perception increases, but as patients' mental pain increases, their pain perception and pain anxiety symptoms decreases (Table 1).

The direct effect of pain perception on the mental pain variable was negative significantly ($P \le 0.001$, Beta = 0.503) and 25% of the variance in mental pain of patients with musculoskeletal pain was explained by their pain perception. Moreover, when pain perception variables (independent variable) and mental pain (mediating variable) were included in the regression equation to predict pain anxiety symptoms in patients with musculoskeletal pain, the effect of pain perception variable on pain anxiety symptoms variable was significantly positive ($P \le 0.001$, Beta =0.435). The effect of mental pain variable on pain anxiety symptoms variable was negative (P=0.015, Beta = -0.154). Finally, both variables (independent and mediating), in total, explain 28% of the variance in pain anxiety symptoms in patients with musculoskeletal pain (Table 2).

The bootstrap test was done to determine the mediating role of mental pain in the relationship between pain perception

Variable	Mean	Standard deviation	Pain perception	Mental pain	Pain anxiety symptoms
Pain perception	52	1.85	1		
Mental pain	157	3.06	-0.05**	1	
Pain anxiety symptoms	171	2.25	0.51**	-0.38**	1

(**) Significance at the level of 0.01

Downloaded from ijmpp.modares.ac.ir on 2024-05-05

Independent	Dependent	Beta	В	Se	Т	Р	R	R ²
Pain perceptio n	Mental pain	-0.50	1.03-	0.11	-9.15	0.001**	0.50	0.25
Pain perception	Pain anxiety symptoms	0.43	0.29	0.04	6.96	0.001**	0.53	0.28
Mental pain		-0.15	-0.05	0.02	-2.46	0.015*		

 Table 2) Regression model for predicting pain anxiety symptoms

(**) Significance at the level of 0.01

(*) Significance at the level of 0.05

Table 3) Bootstrap test results to determine the mediating role of mental pain

Ir va	Independent	Mediating variable	Dependent variable	Resampling	Error estimation	Bootstrap limits			Significance
	variable					Lower limit	Upper limit	if	of P. Value
	Pain perception	Mental pain	Pain anxiety symptoms	1000	0.03	0.01	0.15	0.08	0.04*

(*) Significance at the level of 0.05

and pain anxiety symptoms in patients with musculoskeletal pain. According to Cohen ^[30], the effect size is interpreted as: 0.02(weak), 0.15(medium) and 0.35(strong), respectively. In this study, indirect effect size of pain perception on pain anxiety symptoms in patients with musculoskeletal pain with a standard coefficient of 0.08 at the level of 0.04 was positively significant. As the result, mental pain had a small effect size and it had a mediating role between pain perception and pain anxiety symptoms in patients with musculoskeletal pain. In other words, mental pain causes patients with pain anxiety symptoms to perceive less pain (Table 3).

Discussion

The purpose of this study was predicting pain anxiety symptoms based on pain perception with the mediating role of mental pain in musculoskeletal patients. Results showedmental pain has a mediating role between pain perception and pain anxiety symptoms. This finding is consistent with studies in the research literature. For example, in study of Cimpean and David ^[18] emphasized on the role of pain catastrophizing and state anxiety on pain tolerance and pain-related anxiety. In another view, when cognitive behavioral therapy is conducted on patients, chronic pain, pain anxiety, depression and psychological problems improve ^[19]. In another study, results demonstrated emotional regulation mechanisms in patients with chronic pain are related to their quality of life and life style ^[20].

The intensity of perception of a pain is the intensity of suffering and the concept of pain for the patient. Psychological and cognitive factors such as the ability to understand, pain conceptualization or meaning and personal interpretation are effective in how to respond to pain. It seems pain anxiety is a type of pain perception that is not dependent on external stimuli. On the other hand, recognizing and even measuring pain anxiety can be effective in treating chronic pain. Because pain is expressed in metaphorical statements ^[2], it is often difficult to measure mental and perceived. Often, perceived pain manifests itself in the form of anxiety symptoms pain ^[31]. All three variables of this study are more cognitive and subjective. Therefore, their perceptual and mental aspects are intertwined and they cannot be separated exactly in terms of pain. According to cognitive approach; attentional and conceptual processes direct and manipulate pain and emotional states facilitate our cognitive capacities ^[32].

Although, this study contributed to our knowledge regarding the role of psychological factors in pain perception, it is not without limitations. Due to the online data collecting, it was difficult for patients to cooperate and probabilistic sampling was not possible due to Corona virus outbreak.

Conclusion

Mental pain has a small effect size and it has a mediating role between pain perception and pain anxiety symptoms in patients with musculoskeletal pain. Accordingly, researchers are recommended to pay attention to the mental factors of pain in their future studies.

Acknowledgments

The author hereby announce her gratitude and appreciation to all participants who contribute to this study.

Authors' contribution: RF was corresponding author who helped for drafting introduction and discussion and supervise to design methodology/ and statistical analysis 70%.

RKZ wrote introduction and conduct the research (%20). RR contributed in participants sampling (%10).

Conflicts of Interests: None.

Ethical Permission: In this study, all ethical principles were respected. The purpose of

the study was explained for all participants. Written signed consent form was obtained from all participants

Funding/Support: This research received no financial support.

References

- 1. Raja SN, Carr DB, Cohen M, Finnerup N, Flor H, Gibson S., et al. The revised International Association for the Study of Pain definition of pain: concepts, challenges, and compromises, PAIN, 2020; 161 (9): 1976-1982. doi:10.1097/j. pain.000000000001939.
- 2. Raiisi F. Conceptual Metaphors of Pain in Persian: A Cognitive Analysis. IJMPP. 2021; 6(2): 496-501. http://ijmpp.modares.ac.ir/article-32-50132en.html.
- 3. Raiisi F, Amini F, Khani MM. Investigating the relationship between Skeletal-muscular Pains and Quality of Life among Nursing and Midwifery Students. IJMPP. 2020;5(2): 336-343. Corpus ID: 221755429.
- 4. Amini F, Raiisi F. The Relationship between Musculoskeletal Pain and Demographic Variables in Nursing and Midwifery Students. 2019; 4(4): 256-262.
- 5. Tehranizadeh M., Raiisi F. The Relationships between Depression, Pain Self-Efficacy, Physical Disability and Chronic Pain. IJMPP. 2020;5(3): 373-379. http://ijmpp.modares.ac.ir/article-32-45078-en.html.
- 6. Tossani E. The Concept of Mental Pain. Psychother Psychosom 2013; 82:67-73. doi:10.1159/000343003.
- Mills SEE, Nicolson KP, Smith BH. Chronic pain: a review of its epidemiology and associated factors in population-based studies. Br J Anaesth. 2019, 123(2):e273-e283. doi:10.1016/j.bja.2019.03.023.
- Bushnell M, Čeko M. Low L. Cognitive and emotional control of pain and its disruption in chronic pain. Nat Rev Neurosci, 2013; 14, 502–511. doi:10.1038/nrn3516. doi: 10.1038/ nrn3516.
- 9. Orbach I, Mikulincer M, Sirota P, Gilboa-Schechtman E. Mental pain: a multidimensional operationalization and definition. Suicide Life Threat Behav. 2003;33(3):219-30. doi:10.1521/ suli.33.3.219.23219.
- 10. Verrocchio MC, Carrozzino D, Marchetti D, Andreasson K, Fulcheri M, Bech P. Mental Pain and Suicide: A Systematic Review of the Literature. Front Psychiatry. 2016, 20;7:108. doi:10.3389/ fpsyt.2016.00108.
- 11. Sandler J. Psychology and psychoanalysis. Br J Med Psychol. 1962; 35:91-100.

doi:10.1111/j.2044-8341.1962.tb00507.x.

- 12. Gilam G, Gross JJ, Wager TD, Keefe FJ, Mackey SC. What Is the Relationship between Pain and Emotion? Bridging Constructs and Communities. Neuron. 2020;107(1):17-21. doi:10.1016/j. neuron.2020.05.024.
- 13. Becker S, Navratilova E, Nees F, Van Damme S. Shared Mechanisms of Chronic Pain and Emotional-Motivational Problems: From Basic Science to the Clinics. Pain Res Manag. 2018 19. doi:10.1155/2018/9305026.
- 14. Rajmohan V, Kumar SK. Psychiatric morbidity, pain perception, and functional status of chronic pain patients in palliative care. Indian J Palliat Care. 2013,19(3):146-51. doi:10.4103/0973-1075.121527.
- 15. McGrath PA. Psychological aspects of pain perception. Arch Oral Biol. 1994;39 Suppl:55S-62S. doi:10.1016/0003-9969(94)90189-9.
- Apkarian AV. Pain perception in relation to emotional learning. Curr Opin Neurobiol. 2008, 18(4):464-8. doi:10.1016/j.conb.2008.09.012.
- Peláez I, Martínez-Iñigo D, Barjola P, Cardoso S, Mercado F. Decreased Pain Perception by Unconscious Emotional Pictures. Front Psychol. 2016,21;7:1636.doi:10.3389/fpsyg.2016.01636.
- Cimpean A, David D. The mechanisms of pain tolerance and pain-related anxiety in acute pain. Health Psychology Open. 2019. doi:10.1177/2055102919865161
- 19. Zanini S, Voltolini A, Gragnano G, Fumagalli E, Pagnini F. Changes in Pain Perception following Psychotherapy:TheMediatingRoleofPsychological Components. Pain Res Manag.2018,19; 8713084. doi: 10.1155/2018/8713084.
- 20. Márki G, Bokor A, Rigó J, Rigó A. Physical pain and emotion regulation as the main predictive factors of health-related quality of life in women living with endometriosis. Hum Reprod. 2017; 1;32(7):1432-1438.doi:10.1093/humrep/dex091.
- 21. Hechler T, Endres D, Thorwart A. Why Harmless Sensations Might Hurt in Individuals with Chronic Pain: About Heightened Prediction and Perception of Pain in the Mind. Front Psychol. 2016, 25,7:1638. doi:10.3389/fpsyg.2016.01638.
- 22. Taherizadeh S, Samari A, Ahi G. Structural modeling of pain perception in people with chronic pain syndrome based on behavioral inhibition system: mediated by pain catastrophe and perceived social support. JAP. 2021; 12(2) :19-31. URL:http://jap.

iums.ac.ir/article-1-5574-en.html. [In Persian].

- 23. Melzack R. The McGill Pain properties and scoring methods. Pain, 1975, 1(3), 277-299. doi:10.1016/0304-3959(75)90044-5.
- 24. Dworkin RH, Turk DC, Revicki DA, Harding G, Coyne KS, Peirce-Sandner S, et al. Development and initial validation of an expanded and revised version of the Short-form McGill Pain Questionnaire (SF-MPQ-2). PAIN, 2009, 144(1), 35-42. doi:10.1016/j.pain.2009.02.007.
- Orbach I, Mikulincer M, Sirota P, Gilboa-Schechtman E. Mental pain: A multidimensional operationalization and definition. Suicide and Life-threatening Behavior, 2003; 33(3): 219–230. https://doi.org/10.1521/suli.33.3.219.23219.
- Guimarães R, Fleming M, Cardoso MF. Validation of the Orbach & Mikulincer Mental Pain Scale (OMMP) on a drug addicted population. Soc Psychiatry Psychiatr Epidemiol, 2014; 49, 405– 415. https://doi.org/10.1007/s00127-013-0751-6.
- 27. Guimarães R, Fleming M, Cardoso MF.Validation of the Orbach & Mikulincer Mental Pain Scale (OMMP) on a drug addicted population.Soc Psychiatry Psychiatr Epidemiol. 2014;49(3):405-15.
- 28. karami J, Bagian M, Momeni K, Elahi A. Measurement of mental pain: Psychometric properties and confirmatory factor analysis of multidimensional mental pain questionnaire. Quarterly Journal of Health Psychology, 2018; 7(25): 146-172. [In Persian].
- Paknejad M, Asghari A, Rahiminezhad A, Rostami R, Taheri A. Factorial Structure and Psychometric Properties of the Pain Anxiety Symptoms Scale (PASS-20). Applied Psychological Research Quarterly, 2014, 5(3): 71-94. Doi:10.22059/ JAPR.2014.52307. [In Persian].
- Cohen JE. Statistical Power Analysis for the Behavioral Sciences. Hillsdale, NJ: Lawrence Erlbaum Associates, Inc. 1988.
- 31. CarletonRN,AsmundsonGJ.themultidimensionality of fear of pain: construct independence for the fear of Pain Questionnaire-Short Form and the Pain Anxiety Symptoms Scale-20. J Pain; 2009, 10(1):29-37. doi:10.1016/j.jpain.2008.06.007.
- 32. Pinel L, Perez-Nieto MA, Redondo M, Rodríguez-Rodríguez L, Mateos LL. The Impact of Cognitive Anxiety and the Rating of Pain on Care Processes in a Vigilance Task: The Important Part Played by Age, Pain Research and Management, 2020, https://doi.org/10.1155/2020/3204720.

Downloaded from ijmpp.modares.ac.ir on 2024-05-05